

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

XYLOGLUCAN FUCOSYLTRANSFERASES**FIELD OF THE INVENTION**

This invention is in the field of molecular biology. In particular, this invention relates to the isolation, purification, cloning and expression of plant xyloglucan fucosyltransferases.

This application claims priority from US provisional serial no. 60/117,555 filed January 18, 1999 and US application serial no. 09/490,521 filed January 25, 2000.

BACKGROUND OF THE INVENTION

In most multicellular organisms, cells are embedded in an intricate extracellular matrix that keeps them together and influences the shape, development, and polarity of the cells they contact. Animal cells have such an extracellular matrix at their surface, but plants possess a distinct wall that encloses every cell. Many important differences between plants and animals with respect to nutrition, digestion, growth, reproduction, and defense mechanisms can be traced to the plant cell wall. Cell walls are mediators of growth, which in plants is determined largely by the wall extensibility provided that sufficient turgor pressure is present. Morphogenesis is also effected by the cell wall at the tissue and cellular levels. The biosynthesis of plant cell walls must be very tightly regulated. Although an individual plant cell may expand its volume by as much as 18,840 times, its cell wall must maintain a regular thickness and uniform structure to prevent hemorrhaging of the cell contents due to the high internal turgor pressure. However, despite extensive descriptions of the chemical and physical structure of the plant cell wall, very little is known about its biosynthesis. Only one cell wall-synthesizing glycosyltransferase, cellulose synthase, has been cloned and described in any detail

Plant cell walls are mainly composed of cellulose microfibrils and matrix polysaccharides. Hemicellulose is a type of matrix polysaccharide that binds tightly but noncovalently to cellulose microfibrils, helping to crosslink them into a complex network. Xyloglucan is a fundamentally important hemicellulose in dicot and nongraminaceous monocot plants. It comprises approximately 25% of the total cell wall and forms a load-bearing network by associating with the faces of surrounding cellulose microfibrils via hydrogen bonds. Xyloglucan contains a beta-1,4-glucan backbone decorated with side chains of xylose alone, xylose and galactose, and xylose, galactose and fucose. The presence or absence of the

fucose residue is thought to determine whether the xyloglucan conformation is planar and thus better able to bind to cellulose, a critical step in cell wall formation. In addition, oligosaccharides consisting of a monomer of xyloglucan have been shown to prevent auxin-promoted elongation of pea stems when the

5 oligosaccharides contain fucose, but not if they lack fucose suggesting that xyloglucan fragments act as signalling molecules in vivo. Xyloglucan fucosylation is thus a critical step in plant development.

There is thus a need to identify the genes and gene products involved in plant xyloglucan fucosylation. In particular, there is a need to isolate, purify and

10 clone xyloglucan fucosyltransferase genes and gene products so that xyloglucan fucosylation may be controlled and regulated in plants and other organisms.

SUMMARY OF THE INVENTION

In order to meet these needs, the present invention is directed to purified,

15 isolated, sequenced and cloned plant xyloglucan fucosyltransferase. In addition, the present invention is directed to the purification, isolation, sequencing and cloning of plant xyloglucan fucosyltransferase. The present invention is further directed to transgenic organisms expressing plant xyloglucan fucosyltransferase. The present invention is further directed to transgenic plants expressing regulated

20 levels of xyloglucan fucosyltransferases.

Sub B1 In general, the invention features substantially pure fucosyltransferase DNA or protein obtained from a plant. In a related aspect, the invention features a fragment or analog polypeptide including an amino acid sequence substantially identical to the sequences shown in Tables 2, 5 and 6.

25 *Sub B2* In another related aspect, the invention features substantially pure DNA having a sequence substantially identical to the nucleotide sequence shown in Tables 3-14. In preferred embodiments, such DNA is cDNA or is genomic DNA. In related aspects, the invention also features a vector and a cell (e.g., a plant) which includes such substantially pure DNA. In various preferred embodiments,

30 the vector-containing cell is a prokaryotic cell, for example, E. coli or Agrobacterium or, more preferably, a plant cell.

Sub B3 In yet another related aspect, the invention features a method of fucosylating a polypeptide in vivo involving: (a) providing a cell containing the fucosyl transferase DNA of the invention positioned for expression in the cell; and

35 (b) culturing the transformed cell under conditions for expressing the DNA, resulting in the fucosylation of the protein. In preferred embodiments,

fucosylation occurs in a plant cell.

Sub B4 →

In another aspect, the invention features a recombinant polypeptide fucosylated using a cell expressing DNA which is substantially identical to the nucleotide sequence shown in Tables 3-14. In still other preferred embodiments, the polypeptide is further fucosylated using one or more fucosyl transferases.

Sub B5 →

The present invention further includes multiple types of DNA constructs including (1) "sense" constructs encoding proteins, which can increase the expression of fucosyltransferases in plant species and (2) "antisense" constructs containing DNA, which can be used to produce antisense RNA in to reduce expression of fucosyltransferases in plants. Optimal amounts of antisense RNA in transgenic plants will selectively inhibit the expression of genes in these plants which are involved in the fucosylation of xyloglucans.

Some of these constructs will direct constitutive production of transcripts. Other constructs will direct expression in specific organs and/or specific tissue layers of the transgenic plant. These organs will include leaves, petioles, stems, flower organs, seeds, fruits or photosynthetically active parts of the plant. Tissue layers will include but may not be restricted to the epidermis and adjacent cell layers.

The present invention also provides recombinant cells and plants containing these constructs.

Sub B6 →

In one embodiment, the first category of DNA constructs include: a promoter selected from but not limited to constitutive, tissue-specific, cell-type specific, seed-specific, flower-specific, fruit-specific, epidermis-specific promoters, a promoter specific to cell layers adjacent to the epidermis or a promoter specific to photosynthetically active plant tissues, which functions in plant cells to cause the production of an RNA sequence. In this embodiment, the DNA coding region sequences that encode proteins which can be used to increase the activity of plant fucosyl transferases in transgenic plants. The DNA coding region will further include a region 3' to the coding regions the 3' non-translated region which functions in plant cells to cause the addition of polyadenylate nucleotides to the 3' end of the RNA sequence promoter.

Sub B7 →

In another embodiment, a second category of DNA construct will include a constitutive promoter, seed-specific, flower-specific, fruit-specific, epidermis-specific promoter, a promoter specific to cell layers adjacent to the epidermis or a promoter specific to photosynthetically active plant tissues, which functions in plant cells to cause the production of an RNA sequence. The DNA construct will

also include DNA sequences which can produce antisense RNA molecules. These RNA molecules can selectively inhibit the accumulation of transcripts encoding proteins which encode plant fucosyl transferases.

5 *Sub B8* In accordance with another aspect of the present invention, there is provided a method of producing genetically transformed plants which express a gene or genes involved in fucosyl transferase activity. In this method, a recombinant, double-stranded DNA molecule into the genome of a plant cell. In this embodiment, the DNA sequence will include a promoter which functions in plant cells to cause the production of an RNA sequence in flowers, seeds, fruit or
10 other plant tissues. In addition, the sequence will include a DNA coding sequence encoding proteins involved in fucosyl transferase activity in plants. Alternatively, the sequence will be a template to the synthesis of antisense RNA inhibiting the development of these structures. The DNA sequence will also include a 3' non-translated region which functions in plant cells to cause the addition of
15 polyadenylate nucleotides to the 3' end of the RNA sequences. The method also includes obtaining transformed plant cells and regenerating from the transformed plant cells genetically transformed plants. The transformed plant cells may be used to overproduce in cell culture the fucosylated xyloglucans.

The present invention is also directed to transgenic cells such as yeast,
20 fungi, mammalian, and the like cells expressing the DNA sequences of this invention. The present invention is also directed to purified fucosylated xyloglucans isolated from the transgenic cells of the invention.

Sub B9

BRIEF DESCRIPTION OF THE DRAWINGS

25 This invention will be better understood by reference to the figures, in which:

Figure 1 shows the biochemical purification of xyloglucan-specific fucosyltransferase from pea. Figure 1A shows a silver-stained SDS-PAGE gels showing protein profiles from carbonate-washed pea epicotyl extract, peak
30 fractions from a GDP-agarose and size exclusion column, and the entire profile of all fractions from an anion exchange column. Figure 1B shows a xyloglucan-specific fucosyltransferase activity, in nKat (= nMoles of substrate incorporated into product per second), for each fraction of the anion exchange column eluate.

Figure 2 shows confirmation of Arabidopsis fucosyltransferase (Ftase) activity. Figure 2A shows that Anti-AtFT1 polyclonal antibodies recognize an approximately 62 kDa polypeptide in solubilized membrane proteins of Arabidopsis, but not pea. Lane 1, antigen (50 ng); Lane 2, Arabidopsis (40 :l), Lane 3, pea (40 :l.) Left, Western blot. Right, Commassie staining of membrane showing protein profiles. Figure 2B shows that anti-FTase antibodies immunoprecipitate more XG-specific FTase activity than an equal volume of preimmune serum. Shown is FTase activity, in nanokats, of precipitated pellets that were subsequently subjected to FTase activity assays. This is an example similar to results seen in seven different replicates. Figure 2C shows that a full-length AtFTase which was expressed in a COS cell line shows XG-specific FTase activity. Activity in nKat is shown in the presence or absence of tamarind XG for untransformed Cos-7 cells, cells transformed with vector DNA (pCDNA3), cells transformed with vector containing AtFT1 (pCDNA-XGFT), or solubilized pea Golgi vesicles.

Figure 3 shows a hydrophobicity plot of AtFT1.

Figure 4 shows a diagram of plasmid pMEN020.

DEFINITIONS

To ensure a complete understanding of the invention, the following definitions are provided:

Xyloglucan: . Xyloglucan is a hemicellulose carbohydrate present in dicot and nongraminaceous monocot plants comprising approximately 25% of the total cell wall and forming a load-bearing network by associating with the faces of surrounding cellulose microfibrils via hydrogen bonds. Xyloglucan contains a beta-1,4-glucan backbone decorated with side chains of xylose alone, xylose and galactose, and xylose, galactose and fucose.

Sub B10 **Xyloglucan fucosyltransferase:** Xyloglucan fucosyltransferase (XG Ftase) is an enzyme that fucosylates xyloglucan by adding a fucose residue to xyloglucan.

Transgenic Plants: Transgenic plants are plants which contain DNA sequences which were introduced by transformation.

5 **Promoter:** A promoter is the minimal DNA sequence sufficient to direct transcription. Promoters can render transcription controllable for cell-type specific, tissue-specific, or inducible expression. Promoter elements may be located in the 5' or 3' regions of the native gene.

10 **Poly-A Addition Site:** A poly-A addition site is a nucleotide sequence which causes certain enzymes to cleave mRNA at a specific site and to add a sequence of adenylic acid residues to the 3'-end of the mRNA.

15 **Polypeptide:** Polypeptide means any chain of amino acids, regardless of length or post-translational modification (e.g., glycosylation or phosphorylation).

20 **Substantially Identical:** For a polypeptide, substantially identical means a polypeptide exhibiting at least 50%, preferably 70%, more preferably 90%, and most preferably 95% identity to a reference polypeptide. For a nucleic acid substantially identical means a nucleic acid sequence exhibiting at least 85%, preferably 90%, more preferably 95%, and most preferably 97% identity to a reference nucleic acid sequence. For polypeptides, the length of comparison sequences will generally be at least 16 amino acids, preferably at least 20 amino acids, more preferably at least 25 amino acids, and most preferably 35 amino acids. For nucleic acids, the length of comparison sequences will generally be at least 30 nucleotides, preferably at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 110 nucleotides.

30 Sequence identity is typically measured using sequence analysis software (e.g., Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wis. 53705). Such software matches similar sequences by assigning degrees of homology to various substitutions, deletions, substitutions, and other modifications. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and
35 phenylalanine, tyrosine.

Substantially Pur Polypeptid : Substantially pure polypeptide means a fucosyltransferase polypeptide which has been separated from components which naturally accompany it. Typically, the polypeptide is substantially pure when it is at least 60%, by weight, free from the proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably, the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight fucosyltransferase polypeptide. A substantially pure fucosyltransferase polypeptide may be obtained, for example, by extraction from a natural source (e.g., a plant) by expression of a recombinant nucleic acid encoding a fucosyltransferase polypeptide; or by chemically synthesizing the protein. Purity can be measured by any appropriate method, e.g., column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

A protein is substantially free of naturally associated components when it is separated from those contaminants which accompany it in its natural state. Thus, a protein which is chemically synthesized or produced in a cellular system different from the cell from which it naturally originates will be substantially free from its naturally associated components. Accordingly, substantially pure polypeptides include, without limitation, those derived from eukaryotic organisms but synthesized in *E. coli* or other prokaryotes, or those derived from a eukaryotic cell which does not normally synthesize such a protein, or those derived from a eukaryotic cell engineered to overexpress such a protein.

Substantially Pure DNA : Substantially pure DNA means DNA that is free of the genes which, in the naturally-occurring genome of the organism from which the DNA of the invention is derived, flank the gene. The term therefore includes, for example, a recombinant DNA which is incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of a prokaryote or eukaryote; or which exists as a separate molecule (e.g., a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. It also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequence.

Transformed Cell: Transformed cell means a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a DNA molecule encoding (as used herein) a fucosyltransferase polypeptide.

Positioned for Expression: Positioned for expression means that the DNA molecule is positioned adjacent to a DNA sequence which directs transcription and translation of the sequence (i.e., facilitates the production of, e.g., a recombinant fucosyltransferase polypeptide or RNA molecule).

5

Operably Linked: Operably linked mean that a gene and a regulatory sequence(s) are connected in such a way as to permit gene expression when the appropriate molecules (e.g., transcriptional activator proteins) are bound to the regulatory sequence(s).

10

Purified Antibody: Purified antibody means an antibody which is at least 60%, by weight, free from proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably, the preparation is at least 75%, more preferably 90%, and most preferably at least 99%, by weight, antibody, e.g., a plant fucosyltransferase specific antibody. A purified fucosyltransferase antibody may be obtained, for example, by affinity chromatography using recombinantly-produced fucosyltransferase protein or conserved motif peptides and standard techniques.

15

Specifically Binds: Specifically binds means an antibody which recognizes and binds fucosyltransferase protein but which does not substantially recognize and bind other molecules in a biological sample.

20

DETAILED DESCRIPTION OF THE INVENTION

Taking into account these definitions, the present invention is directed to isolated, purified and cloned plant xyloglucan fucosyltransferases.

25

Plant Fucosyltransferase Purification

Sub B11 → A biochemical approach was utilized to purify sufficient quantities of xyloglucan fucosyltransferases from pea epicotyls. Pea microsomes were prepared as follows: 2 cm segments, excised just below the apical hook, of etiolated *Pisum sativum*, cv Alaska were collected and homogenized in 1.5 volumes buffer (50 mM Hepes pH 7.5, 1 mM EDTA pH 8.0, 0.4 M sucrose, 1 mM DTT, 0.1 mM PMSF, 1 :g/mL each aprotinin, leupeptin, and pepstatin.) The homogenate was filtered, centrifuged at 2,000 x g for 15 minutes, and the supernatant was centrifuged at 100,000 x g for 1 hour. The resulting pellets were washed and homogenized in the presence of 0.1 M Na₂CO₃ to strip away

30

35

peripheral membrane proteins (Y. Fujiki, A. L. Hubbard, S. Fowler, P. B. Lazarow, J. Cell Biol. 93, 97 (1982).) The suspension was centrifuged at 100,000 x g for 1 hour and the resulting pellets were washed and resuspended in buffer (50 mM Pipes-KOH pH 6.2, 20% glycerol, 1 mM EDTA, 1 mM DTT, 0.1 M PMSF, 1 :g/mL each aprotinin, leupeptin, and pepstatin.) The suspension was homogenized, mixed with TritonX-100 to a final volume of 0.8%, and stirred for 1-2 h to solubilize membrane proteins. The suspension was centrifuged a final time at 100,000 x g for 1 h and the supernatant was collected and saved.

Arabidopsis cell suspension culture was also used as a tissue source.

When Arabidopsis cell suspension culture was used, the purification procedure was the same except the cells were lysed in a French pressure cell at 4000 p.s.i.

Pea carbonate-washed supernatants were pooled and separated on a GDP-HA agarose affinity chromatography column and GDP-binding proteins were eluted using excess free GDP. Protein levels were monitored by A280. The protein samples were desalted on a Sephadex G-25 column, concentrated, and further separated on a Phenomenex SEC 4000 size exclusion column. Some samples were further purified using a Poros QE or Resource Q anion exchange column and subsequently column and separated by SDS-PAGE electrophoresis.

20 **Fucosyl Transferase Assay**

A specific assay for fucosyl transferase was developed using tamarind or nasturtium storage xyloglucan, which naturally lack fucosyl residues, as an acceptor and radiolabeled GDP-fucose as a donor [V. Farkas, G. Maclachlan, *Arch. Biochem. Biophys.* **264**, 48 (1988). A. Camirand, D. Brummell, G. Maclachlan, *Plant Physiol.* **84**, 753 (1987)].

Carbohydrate Analysis

To confirm that the purified pea protein synthesizes an alpha-1,2 fucose:galactose linkage, carbohydrate analysis was performed on the product resulting from in vitro fucosylation of tamarind xyloglucan by purified FTase. Carbohydrate linkage analysis of tamarind xyloglucan before (tamarind xg) and after (fucosylated xg) incubation with purified pea FTase. Samples were incubated at room temperature for 20 minutes (for immunoprecipitation samples) or 30 minutes (for protein purification samples) with 25 mM Pipes-KOH pH 6.2, 0.5 mg/mL tamarind xyloglucan, 0.05% ³H GDP-fucose (3.7 mBq/mL, 300 GBq/mM, NEN, Boston, MA). Most assays also contained 50 :M non-radiolabeled

GDP-fucose to provide a quantitative measurement of enzyme activity. Assays of immunoprecipitation samples also contained 5 mM MgCl₂. Reactions were precipitated using 70% ethanol and ³H incorporation was measured by scintillation counting. The amount of fucose incorporated into the product was used to calculate activity in nanokats (nKat = nMoles substrated incorporated into product per second.) The results are shown in Table 1.

Table I

Sugar Residue	% tamarind xg	% Fucosylated xg
4-glucose	16.4	17.5
4,6-glucose	37.0	31.5
t-xylose	19.0	13.5
2-xylose	15.0	14.3
t-galactose	12.6	5.5
2-galactose	-	9.0
t-fucose	-	8.7

10

Tamarind seed xyloglucan was fucosylated by 33 pKat size exclusion column-purified pea FTase (1 mg/mL tamarind XG, 1.5 mM GDP-fucose, 50 mM Pipes-KOH pH 6.2.) XG product was precipitated with ethanol, resuspended in water, reprecipitated, and sent to the Complex Carbohydrate Research Center (Athens, GA) for linkage analysis. An equal amount of tamarind xyloglucan was also submitted for linkage analysis. Linkage analysis indicated that incubation of XG with purified FTase resulted in a decrease in the mole percentage of terminal galactose and the appearance of 2-galactose and terminal fucose, thus verifying the activity of the purified enzyme.

20

Peptide Sequencing

Sub B15 It was possible to purify XG FTase 1400-fold by the end of the size exclusion chromatography step resulting in a total of 0.05 mg protein containing 70 nKat XG FTase activity. After biochemical purification and subsequent assay analysis, two polypeptides of approximately 65 kDa and 60 kDa in size were observed to co-purify repeatedly with XG FTase activity (Figure 1.)

25

Sub B16 Limited peptide sequence was obtained from both proteins. Proteins in size exclusion column eluate fractions containing peak amounts of FTase activity were concentrated using a Millipore 4 mL 10 kDa concentrator and separated by electrophoresis. After brief staining with Coomassie and destaining the separated proteins were excised, rinsed in 50% acetonitrile, stored at -80° C and sent to

30

Harvard Microchemistry (Cambridge, MA) for tryptic peptide sequencing. Six peptide sequences were obtained: VFGFLGR, YLLHPTNNVWGLVVR, AVLITSLSSGYFEK, YYDAYLAK, LLGGLLADGFDEK, and ESILPDVNR.

5 *Sub B17* **Arabidopsis EST Identification**

Sub B18 Using these peptides as a query in the Blastp program identified an Arabidopsis EST, 191A6T7, which contained four out of six peptides in a deduced translation of a potential ORF. The 65 kDa peptide was identified as a homolog of BiP, the usually ER-localized molecular chaperone. It is possible that this
10 chaperone co-purified with FTase activity as an artifact and prevented the denaturation of the FTase during purification, though this has not been confirmed.

Peptides from the lower molecular weight protein were not significantly similar to proteins of known function in databases, but did allow the identification of an Arabidopsis EST which, when translated, contains four out of six peptides
15 with amino acid identity ranging from 63%-85% (See Table 2).

The EST(number 191A6t7) was analyzed to determine if it was a full length clone. Northern blot analysis using the ~900 bp-long 191A6T7 as a probe detected an approximately 2 kb transcript, indicating that the EST did not contain the full-length cDNA (RMP, data not shown.).

20 191A6T7 was used as a probe to screen the CD4-15 portion of a size-fractionated Arabidopsis cDNA library at high stringency (J - J - Kieber, M. Rothenberg, G. Roman, K. A. Feldmann, J. R. Ecker, *Cell* **72**, 427 (1993). Two cDNA clones were isolated, the longest containing a 1768 bp insert. Both lacked 13 nucleotides of the 3' UTR and the poly-A tail found in 191A6T7. There
25 is an AATAAA consensus polyadenylation signal eight nucleotides from the 3' end of the library-derived clones. The sequence contains a 1698 nucleotide ORF that encodes a 63.7 kDa protein a 1698 nt open reading frame (Table 3) and correspond to a region of the fully sequenced Arabidopsis bacterial artificial chromosome (BAC) T18E14 (Table 4).

30 The cDNA and corresponding genomic clone have been designated AtFT1. Interestingly, analysis of the BAC indicates that there may be a second FTase approximately 600 bp downstream from AtFT1 which is ~60% identical to AtFT1 (Table 5). Whether this second FTase is expressed, as well as splicing patterns and localization of the encoded protein, are matters of current
35 investigation. It does raise the possibility that a multi-gene family of FTases may exist in Arabidopsis. We will determine whether members of such a family might

be differentially regulated by such factors as environmental stress, tissue localization, or developmental stage. Alternatively, there may well be FTases which have different acceptors, such as carbohydrate protein modifications.

5 **Antibody Preparation**

In order to confirm the identity of AtFT1 as encoding a xyloglucan-specific fucosyltransferase, we prepared polyclonal antibodies directed against AtFT1 and used them to immunoprecipitate proteins from carbonate-washed, solubilized Arabidopsis proteins.

10 *Sub B19* The portion of AtFT1 encoding aa 73 to 566 was PCR-amplified using appropriate primers and cloned into the pET28a expression vector (Novagen, Madison WI.) The resulting insoluble fusion protein was purified by washing four times with 1% Triton X-100, 50 mM Hepes-KOH pH 7.6, 10 mM MgCl₂ and one time with 25 mM Hepes-KOH pH 7.0, 8 M urea. The pellet was resuspended in 6
15 M guanidine-HCl and protein was precipitated from the supernatant with 10% TCA. The protein was emulsified with Titermax adjuvant (CytRx Corporation, Norcross, GA) and injected into a rabbit. For western blotting, 40 :l of carbonate-washed solubilized protein from pea and Arabidopsis and 50 ng of purified antigen were separated by SDS-PAGE and electroblotted. Anti-AtFT1
20 Abs (1:5000) were used for western blotting. Goat-antirabbit antibodies conjugated to horseradish peroxidase was used as a secondary antibody. Signals were detected by the enhanced chemiluminescence method (Pierce, Rockford, IL.) Membranes were then stained with Coomassie blue to detect protein.

25 **Immunoprecipitations**

Sub B20 For immunoprecipitations, solid NaCl was added to carbonate-washed solubilized Arabidopsis protein to a final concentration of 200 mM. The Arabidopsis protein was precleared by incubation with 1/10 volume of 50% slurry of protein A sepharose beads (Pharmacia) in buffer A (25 mM Pipes-KOH pH 7.5,
30 50 mM NaCl, 2 mM EDTA pH 8.0.) The resulting supernatants were incubated with 50 :l of immune or preimmune anti-AtFT1 serum for 1 h. 1/5 volume of protein A sepharose slurry was added to precipitate the antigen-antibody complexes and the samples were incubated for an additional 3 hours with rocking at 4 degrees C. Samples were then centrifuged, washed five times in buffer A containing 1%
35 Triton X-100 and two times in buffer A without detergent. The pellets were

resuspended in buffer A to a final volume of 120 :l and assayed for AtFTase activity as described above.

5 The immunoprecipitated proteins were then assayed for XG FTase activity. More FTase activity was correlated with pellets derived from immunoprecipitation reactions using immune antiserum rather than preimmune serum, thereby indicating that the Arabidopsis clone encodes a xyloglucan-specific FTase (Figure 2).

Expression in COS Cells

10 *Sub B22* Cos-7 cells were grown on 100 mm plates in DMEM-10% Fetal Bovine Serum. Cells were transfected with different plasmids using Lipofectamine™ reagent (Life Technologies) following the manufacturer's instructions using 9 :g of DNA and 72 :g of Lipofectamine. Cells were incubated for 24 hours in the medium containing DNA-Lipofectamine without Fetal Bovine Serum. The medium
15 was changed to DMEM-10% Fetal Bovine Serum and incubated for another 48 hours. The cells were scraped off the dish in 0.25 M sucrose, 10 mM Tris-HCl pH 7.5 and 0.4% CHAPS. XG-FTase activity was measured using 50 :g of protein in the absence (-XG) or presence (+XG) of 100 :g tamarind xyloglucan. The incubation was carried out in a volume of 0.1 mL in the presence of 1 :M GDP-Fuc;
20 (93,000 dpm), 10 mM MnCl₂, 20 mM Hepes pH 7.0, 0.05% Triton X-100 at 25°C for 90 min. The reaction was halted by adding ethanol to a final concentration of 70%. Samples were incubated at 4°C and filtered through 1.5 :m glass fiber filters. The filters were washed with 70% ethanol containing 1 mM EDTA. The filters were dried and radioactivity determined by liquid scintillation. A control
25 using pea Golgi vesicles was carried out in parallel. The results indicate that AtFT1 expressed in a COS cell line showed in vitro FTase activity that was 41 times higher than COS cells transformed with an empty vector and 1.4 times higher than solubilized pea Golgi vesicles (Figure 2).

30 Taken together, the in vitro translation data and the cos cell translation data provide strong evidence that AtFT1 is involved in xyloglucan biosynthesis.

Sequence Analysis of AtFT1

35 *Sub B23* Analysis of AtFT1 indicates that, while it has some structural characteristics common to other fucosyltransferases, it is quite divergent at the amino acid sequence level. Hydrophobicity plots predict that there may be a

N-terminal transmembrane signal anchor sequence (see Figure 3). In vitro translation in the presence of canine pancreatic microsomes followed by carbonate washing of the products indicates that the AtFT1 translation product is a membrane protein (data not shown.) As with other glycosyltransferases, the C-terminal region is predicted to be largely hydrophilic.

AtFT1 is not significantly similar to any other FTases from other organisms, although multiple sequence alignments have identified three motifs which appear to be conserved among all alpha 1,2-FTases. One ([IV]G[IV]HQ)[VI]R..[DN]) has been described previously (Breton et al., 1998). In addition, a second motif (D[EK]..F.[EQ].DQ) and a third hydrophobic region was conserved.

Since these proteins have different acceptor molecules but share the same sugar nucleotide donor (GDP-fucose), it is possible that these regions are involved in GDP-fucose binding or that are necessary for assumption of conserved structural characteristics. Some small regions of similarity are observed between AtFT1 and NodZ, a fucosyltransferase in *Rhizobium* involved in the synthesis of nodulation factors.

Other Glycosyltransferases

The unique nature of this FTase allow its use as a tool for identifying other glycosyltransferases. Consideration of the number of different linkages present in plant cell wall polysaccharides indicates that there should be several hundred different glycosyltransferases involved in cell wall biosynthesis. Several other sequences in the Arabidopsis databases do appear to be similar to AtFT1 and AtFT2, and thus might represent a multi-gene family of FTases or glycosyltransferases. Substantially identical sequences are presented in Tables 6-14.

In addition to database analysis, the isolation of additional plant fucosyltransferases is made possible using standard molecular biology techniques. In particular, using all or a portion of the amino acid sequence of a plant fucosyltransferase of the invention, one may readily design fucosyltransferase oligonucleotide probes, including fucosyltransferase degenerate oligonucleotide probes (i.e., a mixture of all possible coding sequences for a given amino acid sequence). These oligonucleotides may be based upon the sequence of either strand of the DNA comprising the motif. General methods for designing and preparing such probes are provided, for example, in Ausubel et al., and Guide to Molecular Cloning Techniques, 1987, S.

L. Berger and A. R. Kimmel, eds., Academic Press, New York. These oligonucleotides are useful for fucosyltransferase gene isolation, either through their use as probes capable of hybridizing to fucosyltransferase complementary sequences or as primers for various polymerase chain reaction (PCR) cloning strategies. In one particular example, isolation of other fucosyltransferase genes is performed by PCR amplification techniques well known to those skilled in the art of molecular biology using oligonucleotide primers designed to amplify only sequences flanked by the oligonucleotides in genes having sequence identity to fucosyltransferase of the invention. The primers are optionally designed to allow cloning of the amplified product into a suitable vector.

Hybridization techniques and procedures are well known to those skilled in the art and are described, for example, in Ausubel et al. If desired, a combination of different oligonucleotide probes may be used for the screening of the recombinant DNA library. The oligonucleotides are labelled with ³²P using methods known in the art, and the detectably-labelled oligonucleotides are used to probe filter replicas from a recombinant DNA library. Recombinant DNA libraries may be prepared according to methods well known in the art, for example, as described in Ausubel et al., supra, or may be obtained from commercial sources.

For detection or isolation of closely related fucosyltransferases, high stringency conditions may be used; such conditions include hybridization at about 42 degrees C. and about 50% formamide; a first wash at about 65 degrees C., about 2X SSC, and 1% SDS; followed by a second wash at about 65 degrees C. and about 0.1% SDS, 1X SSC. Lower stringency conditions for detecting fucosyltransferase genes having about 85% sequence identity to the fucosyltransferase gene described herein include, for example, hybridization at about 42 degree C in the absence of formamide; a first wash at about 42 degrees C, about 6X SSC, and about 1% SDS; and a second wash at about 50 degrees C., about 6X SSC, and about 1% SDS.

Fucosyltransferase oligonucleotides may also be used as primers in PCR cloning strategies. Such PCR methods are well known in the art and described, for example, in PCR Technology, H. A. Erlich, ed., Stockton Press, London, 1989; PCR Protocols: A Guide to Methods and Applications, M. A. Innis, D. H. Gelfand, J. J. Sninsky, and T. J. White, eds., Academic Press, Inc., New York, 1990; and Ausubel et al., supra. If desired, fucosyltransferases may be isolated using the PCR "RACE " technique, or Rapid Amplification of cDNA Ends (see, e.g., Innis et al., supra). By this method, oligonucleotide primers based on a fucosyltransferase conserved domain are oriented in the 3' and 5' directions and are used to

generate overlapping PCR fragments. These overlapping 3'-and 5'-end RACE products are combined to produce an intact full-length cDNA. P.N.A.S. 85:8998(1988)

5 **Plant Transformation**

Sub B2 Once identified, fucosyl transferase genes can be expressed in a variety of cells including plant cells, yeasts, fungi, bacterial cells and mammalian cells. A wide variety of plants can be transformed to express fucosyl transferase genes and genes related to fucosyl transferase in order to regulate plant carbohydrate glycosylation.

A. Dicots

Methods for transforming a wide variety of different dicots and obtaining transgenic plants are well documented in the literature (See Gasser and Fraley 15 (1989) *Science* 244:1293; Fisk and Dandekar (1993) *Scientia Horticulturae* 55:5-36; Christou (1994) *Agro Food Industry Hi Tech* (March/April 1994) p.17, and the references cited therein).

B. Monocots

Methods for producing transgenic plants among the monocots are currently available. Successful transformation and plant regeneration have been achieved in asparagus (*Asparagus officinalis*; Bytebier et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:5345); barley (*Hordeum vulgare*; Wan and Lemaux (1994) *Plant Physiol* 104:37); maize (*Zea mays*; Gordon-Kamm et al., (1990) *Plant Cell* 2:603; Fromm et al. (1990) *Bio/Technology* 8:833; Koziel et al. (1993) *Bio/Technology* 11:194); oats (*Avena sativa*, Somers et al. (1992) *Bio/Technology* 10:1589); orchardgrass (*Dactylis glomerata*; Horn et al. (1988) *Plant Cell Rep.* 7:469); rice (*Oryza sativa*, including indica and japonica varieties; Toriyama et al. (1988) *Bio/Technology* 6:10; Zhang et al. (1988) *Plant Cell Rep.* 7:379; Luo and Wu (1988) *Plant Mol. Biol. Rep.* 6:165; Zhang and Wu (1988) *Theor. Appl. Genet.* 76:835; Christou et al. (1991) *Bio/Technology* 9:957; rye (*Secale cereale*; De la Pena et al. (1987) *Nature* 325:274); sorghum (*Sorghum bicolor*; Cassas et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:11212); sugar cane (*Saccharum spp.*; Bower and Birch (1992) *Plant J.* 2:409); tall fescue (*Festuca arundinacea*; Wang et al. (1992) *Bio/Technology* 10:691); turfgrass (*Agrostis palustris*; Zhong et al. (1993) *Plant Cell Rep.* 13:1); wheat (*Triticum aestivum*; Vasil et al. (1992)

Bio/Technology 10:667; Troy Weeks et al. (1993) *Plant Physiol.* 102:1077; Becker et al. (1994) *Plant J.* 5:299).

C. Expression Vectors

5 *Substantive*
A variety of expression vectors can be used to transfer the gene encoding plant fucosyl transferase activity as well as the desired promoters and regulatory proteins into a plant. Examples include but not limited to those derived from a Ti plasmid of *Agrobacterium tumefaciens*, as well as those disclosed by Herrera-Estrella, L., et al., *Nature* **303**: 209 (1983), Bevan, M., *Nucl. Acids Res.* **12**: 8711-8721 (1984), Klee, H. J., *Bio/Technology* **3**: 637-642 (1985), and EPO Publication 10 120,516 (Schilperoort et al.) for dicotyledonous plants. Alternatively, non-Ti vectors can be used to transfer the DNA constructs of this invention into monotyledonous plants and plant cells by using free DNA delivery techniques. Such methods may involve, for example, the use of liposomes, electroporation, 15 microprojectile bombardment, silicon carbide whiskers, viruses and pollen. By using these methods transgenic plants such as wheat, rice (Christou, P., *Bio/Technology* **9**: 957-962 (1991)) and corn (Gordon-Kamm, W., *Plant Cell* **2**: 603-618 (1990)) are produced. An immature embryo can also be a good target tissue for monocots for direct DNA delivery techniques by using the particle gun 20 (Weeks, T. et al., *Plant Physiol.* **102**: 1077-1084 (1993); Vasil, V., *Bio/Technology* **10**: 667-674 (1993); Wan, Y. and Lemeaux, P., *Plant Physiol.* **104**: 37-48 (1994), and for *Agrobacterium*-mediated DNA transfer (Hiei et al., *Plant J.* **6**: 271-282 (1994); Rashid et al., *Plant Cell Rep.* **15**: 727-730 (1996); Dong, J., et al., *Mol. Breeding* **2**: 267-276 (1996); Aldemita, R. and Hodges, T., *Planta* **199**: 612-617 (1996); Ishida et al., *Nature Biotech.* **14**: 745-750 (1996)). In addition, plasmid 25 pMEN020 is described in Figure 4.

D. Plant Regeneration

After transformation of cells or protoplasts, the choice of methods for 30 regenerating fertile plants is not particularly important. Suitable protocols are available for *Leguminosae* (alfalfa, soybean, clover, etc.), *Umbelliferae* (Carrot, celery, parsnip), *Cruciferae* (cabbage, radish, rapeseed, broccoli, etc.), *Curcubitaceae* (melons and cucumber), *Gramineae* (wheat, corn, rice, barley, millet, etc.), *Solanaceae* (potato, tomato, tobacco, peppers, etc.), and various 35 other crops See protocols described in Ammirato et al. (1984) *Handbook of Plant Cell Culture -Crop Species*. Macmillan Publ. Co. Shimamoto et al. (1989)

Nature 338:274-276;; Fromm et al. (1990) *Bio/Technology* 8:833-839. ; Vasil et al. (1990) *Bio/Technology* 8:429-434.

E. Carbohydrates from Transgenic Plants

5 Once transgenic plants are produced, carbohydrates can be isolated from the plants by procedures well known in the art. These purified carbohydrates are useful in agriculture as well medicine.

F. New Complex Carbohydrates

10 The enzymes involved in xyloglucan biosynthesis are reasonably stable and moderately abundant in plants. As such, these enzymes find use in synthesizing various types of complex carbohydrates under controlled conditions. It is also possible to make new complex carbohydrates that do not exist in Nature by procedures well known in the art.

G. Carbohydrates as Herbicides

15 *Sub B26* Fucoxyloglucan (XG) is the major hemicellulosic polysaccharide in the primary cell wall of dicots. Monocots have small quantities of XG, but it seems to be much less important in all monocots including grasses. XG has a backbone of
20 beta-1,4 linked glucosyl residues, with three out of every four residues substituted with xylose in a regular repeat, i.e. three substituted followed by one free. Approximately one out of six of the xylosyl residues is further substituted with galactose and on the two position of galactose is an alpha linked fucosyl residue. Thus, the fucose is a peripheral sugar in this polymer.
25 However, the fucose has been postulated to be very important in determining the conformation of the polysaccharide, including controlling interactions of XG with cellulose. Thus, the presence of fucose may be important for the function of this polysaccharide.

Sub B27
30 Since XG is the major hemicellulosic polysaccharides in dicots, including many important weeds, but not very abundant in the cell walls of monocots, including corn, wheat, rice, barley, etc., inhibitors of XG synthesis may be valuable herbicides. Inhibitors include specific inhibitors of the enzyme itself and antisense constructs for inhibiting expression of the protein.

It appears that all of the enzymes that synthesize XG are part of a complex
35 thereby permitting the use of XG-specific fucosyltransferase to identify other enzymes involved in XG synthesis. With the complete set of XG biosynthetic

enzymes in hand, rationale herbicide design is feasible by procedures well known in the art.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236

Table 2

Deduced Amino Acid Sequence from the cDNA

5 MDQNSYRRRSSPIRTTTGGSKSVNFSELLQMKYLSSGT
MKLTRTFTTCLIVFSVLVAFSMIFHQHPSDSNRMGFAEA
RVLDAGVFPNVTNINSDKLLGGLLASGFDEDSCLSRYS
VHYRKPSYPKSSYLISKLRNYEKLHKRCGPGTESYKKA
LKQLDQEHIDGDGECKYVWVWISFSLGNRILSLASVFLYA
10 LLTDRVLLVDRGKDMDDLFCPEFLGMSWLLPLDFPMTD
QFDGLNQESSRCYGYMVKNQVIDTEGTLHLVHLD
YGDHDKMFFCEGDQTFIGKVPWLIVKTDNYFVPSLWLIP
GFDDELNKLFPQKATVFHHLGRYLFHPTNQVWGLVTRY
YEAYLSHADEKIGIQVRVFDEDPGPFQHVMDQISSCTQK
15 EKLLPEVDTLVERSRLVNTPKHKAVLVTSNAGYAENLK
SMYWEYPTSTGEIIGVHQPSQEGYQQTEKKMHNGKALA
EMYLLSLTDNLVTSWSTFGYVAQGLGLKPWILYRPEN
RTTPDPSCGRAMSMEPCFHSPPFYDCKAKTGIDTGLV
20 PHVRHCEDISWGLKLV

B

Table 3
cDNA Sequence

5 ATGGATCAGAATTCGTACAGGAGAAGATCGTCTCCGAT
 CAGAACCACTACCGGCGGTTCAAAGTCCGTTAATTTCTC
 CGAACTACTTCAAATGAAGTATCTCAGCTCCGGTACGAT
 GAAGCTCACGAGAACCTTCACTACTTGCTTGATAGTCTT
 CTCTGTACTAGTAGCATTCTCAATGATCTTTCACCAACA
 CCCATCTGATTCAAATCGGATTATGGGTTTCGCCGAAGC
 10 TAGAGTTCTCGACGCCGGAGTTTTCCCAAATTCTGATAA
 GCTTCTCGGAGGGCTACTTGCTTCTGGTTTTGATGAAGA
 TTCTTGCCTTAGTAGGTACCAATCAGTTCATTACCGTAA
 ACCTTCACCTTACAAGCCATCTTCTTATCTCATCTCTAAG
 CTTAGAAACTACGAAAAGCTTCACAAGCGATGTGGTCC
 15 GGGTACTGAATCTTACAAGAAAGCTCTAAAACAACCTTGA
 TCAAGAACATATTGATGGTGATGGTGAATGCAAATATGT
 TGTGTGGATTTCTTTTAGCGGCTTAGGGAACAGGATACT
 TTCTCTAGCCTCGGTTTTTCTTACGCGCTTTTAACGGAT
 AGAGTCTTGCTTGTGACCGAGGGAAAGACATGGATGA
 20 TCTCTTTTGCAGCGCTTTCTCGGTATGTCGTGGTTGCT
 ACCTTTAGATTTCCCTATGACTGATCAGTTTGATGGATTA
 AATCAAGAATCATCTCGTTGTTATGGATATATGGTGAAG
 AATCAGGTGATTGATACTGAGGGAACCTTGTCTCATCTT
 TATCTTCATCTTGTTCATGATTATGGAGATCATGATAAGA
 25 TGTCTTCTGTGAAGGAGACCAACATTCATCGGGAAAG
 TCCCTTGGTTGATTGTTAAAACAGACAATTACTTTGTTCC
 ATCTCTGTGGTTAATACCGGGTTTCGATGATGAACTAAA
 CAAGCTATTCCACAGAAAGCGACTGTCTTTCATCACTT
 AGGTAGGTATCTTTTTACCCAACTAACCAAGTATGGGG
 30 CTTAGTCACTAGATACTACGAAGCTTACTTATCGCATGC
 GGATGAGAAGATTGGGATTCAAGTAAGAGTTTTCGATGA
 AGACCCGGGTCCATTTACGCATGTGATGGATCAGATTTC
 ATCTTGTA CTCAAAAAGAGAACTTCTACCTGAAGTAGA
 CACACTAGTGAGAGATCTCGCCATGTTAATACCCCAA
 35 ACACAAAGCCGTGCTTGTCACATCTTTGAACGCGGGTTA
 CGCGGAGAACTTAAAGAGTATGTATTGGGAATATCCGA
 CATCAACTGGAGAAATCATCGGTGTTATCAGCCGAGC
 CAAGAAGGTTATCAGCAGACCGAAAAAAGATGCATAAT
 GGCAAAGCTCTTGCGGAAATGTATCTTTGAGTTTGACA
 40 GATAATCTTGTGACAAGTGCTTGGTCTACATTTGGATAT
 GTAGCTCAAGGTCTTGGAGGTTTAAAGCCTTGGATACTC
 TATAGACCCGAAAACCGTACAACCTCCCGATCCTTCGTGT
 GGTGCGGGCTATGTCGATGGAGCCTTGTTCCTCACTCGCC
 TCCATTCTATGATTGTAAAGCGAAAACGGGTATTGACAC
 45 GGGAACACTAGTTCCTCATGTGAGACATTGTGAGGATAT
 CAGCTGGGGACTTAAGCTAGT ATGA

Table 4
Genomic Sequence

5 **BAC T18E12 Accession AC005313 nucleotides 41227-43179 (contains an intron).**

atggatcagaattcgtacaggagaagatcgtctccgatcagaaccactaccggcgggtca
 10 aagtcgtaatttctccgaactacttcaaataagatctcagctccggtacgatgaag
 ctacagagaaccttactactgttgatagcttctctgtactagtagcatttcaatg
 atctttaccaacacccatctgattcaaatcggattatgggttcgccgaagctagagt
 ctgacgccggagtttcccaaatgttactaacatcagatgtgttctccaagtcaaag
 tttgagctttattactttagatctcgttcttactacacgcatttgcctctgtatgcc
 15 atagctcttggtcgatttcaattgagatctatactcataaaaaatgagctttgtcagt
 cacaagactactattttgggttgatgtgttttggtaaaaagtgtctttgtttgg
 tctcagcttagactgttacattcgtttttccgagtttttagatttgttctgattctg
 tttgtttgtagattctgataagcttctcggagggtactgtcttctggtttgatgaa
 gattcttgccttagtaggtaccaatcagttcattaccgtaaaccttcacctacaagcca
 20 tcttctatctcatcttaagcttagaaactacgaaaagcttcacaagcgatgtggtccg
 ggtactgaatcttacaagaaagctctaaaaaacttgatcaagaacataattgatggtgat
 ggtgaatgcaaatagttgtgtgtgatttcttttagcggcttagggaacaggatacttct
 ctagcctcggttttcttaccgcgttttaacggatagagtctgtgttgaccgaggg
 aaagacatggatgatctcttttgcgagccgttctcggatgtcgtgtgtgtaccttta
 25 gatttccctatgactgatcagtttgatggattaaatcaagaatcatctcgtgttatgga
 tatatgggaagaatcagggtattgatactgagggaaacttgtctcatcttattctcat
 ctgttcatgattatggagatcatgataagatgttcttctgtgaaggagaccaaacttc
 atcgggaaagtcccttgggttattgttaaaacagacaattacttgttccatctctgtgg
 ttaataccgggttctgatgatgaactaaacaagctatcccacagaaagcgactgtctt
 30 catcacttaggttaggtatcttttaccaccaactaaccaagtatggggcttagtcactaga
 tactacgaagcttacttaccgatgcggtatgagaagattgggattcaagtaagagtttct
 gatgaagaccgggtccatttcagcatgtgatggatcagatttcatctgttactcaaaaa
 gagaacttctacctgaagtagacacactagtggagagatctcgccatgttaataccccc
 aaacacaaagccgtgctgtcacatcttgaacgcgggttacgcggagaacttaagagt
 atgtattgggaatatccgacatcaactggagaaatcatcgggttcatcagccgagccaa
 35 gaagggtatcagcagaccgaaaaaaagatgcataatggcaaagcttgcggaaatgtat
 cttttgagtttgacagataatctgtgacaagtgttggctacatttgatattgtagct
 caaggcttggagggttaagccttggatactctatagaccgaaaaccgtacaactccc
 gatccttctgtgtgtcgggctatgtcgtatggagccttgttccactgcctccattctat
 gattgtaaagcgaacgggtattgacacgggaacactagttcctcatgtgagacattgt
 40 gaggatatcagctggggacttaagctagtatga

Table 5

Adjacent h m logous g ne on BAC

BAC T18E12 Acc ssion AC005313 nucle tides 43562-45245

5 atgagaatcacagagatcttagcttggttcatggtttagtccctgtctcgctagtaatc
 gtagccatggttgatatgatcaaggaaatggcttgtacaagcatctagattcataaca
 atggaaccaaattgacatcctcatcagatgattcatcactagtcagagagatcaagaa
 caaaaaggtaaaacttacttctcttttggaaatgttctaaattttcttgaa
 tgtttcatcagattctgtagatatgtctctgctggagggtactgtatctggttcaa
 10 gaaagagcttgccttgagtagataccaattaccttaccgttaaagcttcaccgtataa
 acctcgttgcatctacttctgaagcttagagcttacgaagagcttcataaaagatgtgg
 accgggaacaagacagatataccaatgcagaaagattgcttaaacagaaacaaacaggtga
 gatggaatcacaaggatgaagatgtgttggatgtcgtttagcggattaggaacag
 gattatcagattgtctgtgttctgtatgcaatgttgacagatagagcttgcctgt
 15 tgaaggaggggaacagttcgcggatttattctgcgaaccgttctcgataccacttgggt
 actaccgaaagattcaccttagctagtcagttcagttggcttgggtcaaaactcagctca
 ctgcatgagagatgtctgaaggaggaaactgattaatgaatcctctgttctgtctgtc
 tcatctctatctcatctagctcatgactacaatgagcacgacaaaatgttcttctgtga
 agaagatcaaaatctctaaagaatgttccctgggtgatcatgaggacaaacaacttct
 20 tgcaccgtctcttcttatttcttcttctgaagaagagctcgggtatgatgttcccga
 gaaaggaacgggtttaccatttaggtcgttacctttccatccttcaaatcaagctcg
 gggactaatcacaagatactatcaagcttacttagccaaagctgatgaaaggattgggtct
 tcaataagagctttagtgagaaatccggcgtatctctcgagtcacaaagcaaatcat
 ttcgtgtgtcaaaacgagaatctgttaccgagactaagcaaaaggtgaagaacaatacaa
 25 gcagccatcagaagaagagttgaaactcaaatctgttgggtcacctcttaacaacagg
 atactttgagatcttgaaaacaatgtattgggaaaatccaactgtaacaagagatgtgat
 tggaaatcatcagccaagtcagtaaggacatcaacaacagagaagctaatagcataacag
 gaaagcttgggcagagatgtacttactcagcttaacggataagttgggtattagtgctg
 gtctacatttgggtatgtagctcaaggacttgaggattaagagcttggattctgtataa
 30 acaagagaatcaaaccaacccaaatccaccttgcggtagagctatgtcaccagatccttg
 ttccatgtctctcttactatgattgcaaagcaaagaaaggaactgacactggtaattgt
 tgtccgcagtgtagacattgtgaagatatagctggggacttaagcttgggtgacaact
 tttag

35 **Protein translation:**

MRITEILALFMVLVPVSLVIVAMFGYDQGNFVQASRFITMEPN
 VTSSDDSSLVQRDQEQKDSVDMSLLGGLLVSGFKKESCLSRYSYLYRKASP
 YKPSLLLSKLRAYEELHKRCGPGRQYTNAERLLKQKQTGEMESQGCKYVWWM
 40 SFSGLGNRISIASFVLYAMLTDRVLLVEGGEQFADLFCEPFLDTTWLLPKDFTLA
 SQFSGFGQNSAHCHGDM LKRKLINSSVSSLSHLYLHLAHDYNEHDKMFFCEE
 DQNLLKNVPWLIMRTNFFAPSLFLISSFEEELGMMFPEKGTVFHHLGRYLFHPS
 NQVWGLITRYYQAYLAKADERIGLQIRVFDEKSGVSPRVTKQIISCVQENLLPRL
 SKGEEQYKQPSEELKLKSVLVTSLTTGYFEILKTMWENPTVTRDVIGIHQPSH
 45 EGHQQTEKLMHNRKAWAEMYLLSLTDKLVISAWSTFGYVAQGLGLRAWILYK
 QENQTNPNPPCGRAMSPDPCFHAPPYYDCKAKKGTDTGNVVPVHRHCEDISW
 GLKLVDNF

Tabl 6

Arabidopsis thaliana cDNA clone 202C15T7, mRNA s quence [Arabidopsis thaliana]

5

DNA Sequence

TGTTCCATCNTTATGGTTTAATCCAACNTCCAAACCGAACTAACGAAGCTGT
TTCCGCANAAAGAAACCGTGTTTCACCACTTGGGTCGGNATCTTTTACCCT
AAAAATCAAGTTTGGGATATCGTCACNAAGTACTACCATGNTCACTTATCCAA
AGCAGATGNGAGACTCGGGATTCAAATTCGGGTTTTTNGCGATCAAGGTGGA
TACNACCAACACGTCATGGACCAGGTCATATCCTGCACACA

10

Translation of 202C15T7 in correct ORF.

15

VPSLWFNPTXQTELTCLFPXKETVFHHLGRXLFXPKNQVWDIVTKYYHXHLSKA
DXRLGI QIRVFXDQGGYXQHVMQVISCT

Tabl 7

Arabidopsis thaliana cDNA clone 170K19T7, mRNA sequence [Arabidopsis thaliana]

5 TGGNATTACAGATTACAAAGATACGAGGNTCTTCATAGACGTTGTGGACCATT
CACTAGATCCTATAACTTAACACTTGACAACTCAAGTCGGGAGATCGGTCTG
ACGGTGAAGTTTCTGGTTGTAGATATGTAATATGGTTGANTTCCAATGGTGAT
10 CTTGGGAATAGGATGCTGAGTCTAGCTTCANCTTTNCTTTATGCTCTCTTAAC
AAATAGGTTTTNACTTGTCGAACTAGGAGTTGACATGGCTGATCTTTTCTNCG
AGCCATTTCCAAACACTACTTGTTTTCTCCCCCAGAGTTTCCGCTCAACAGC
CACTTCAACGAGCAAGTCTCTTTCTAACGGAAATTNTTGGCAACCCCGATGG
15 GTTCATAATCGNNCATGTAGTTCCGTNATTCCCAGTGNCCAACAAAAAGCTTT
TTNTTTTTGNNAGGNTAGCCAAGTTTTTTTNGGGGAAACCCCTGGTTGTCTT
AAAANCGGGTAGNT TTTTTTCCCAACTTTTTTTNA

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171

Table 8
T31J1TR TAMU Arabidopsis thaliana gen mic clon T31J1, genomic survey
sequ nce

5 CAAGCTTACAAGAAAGCAACGGAGATTCTTGGTCATGATGATGAGAATCATTC
AACCAAATCTGTTGGTGAATGCAGATACATTGTGTGGATTGCTGTTTATGGGC
TAGGAAACAGAATACTTACTCTTGCTTCTCTGTTTCTCTATGCTCTCTTGACTG
ACAGAATCATGCTTGTTGACCAACGTACGGACATAAGTGACCTCTTCTGTGA
GCCTTTTCCAGGTACTTCCTGGCTACTCCCTCTGGATTTTCCACTAACAGATC
10 AATTAGATAGCTTCAACAAGGAATCTCCGCGCTGTTACGGAACAATGTTGAA
GAATCATGCCATTAACCACTACAACAGAAAGCATCATCCCCTCGTACCTCT
GTCTTTATCTTATTCACGATTACGACGATTATGATAAGATGTTCTTCTGTGAAA
GTGACCAAATTCTCATCAGGCAAGTCCCTTGGTTGGTCTTCAACTCGAATCTT
TACTTTATCCCATCTCTATGGTTGATCCCTTCTTTTCAGTCAGAATTAAGCAAG
15 CTATTCCCACAGAAAGAAACCGTCTTTCACCATTTGGCTCGCTATCTTTTCA
CCCGACTAACCAAGTTTGGGGCATGATCACAAGATCCTATAATGGGTATTTAT
CAAGAGCTGATGAGAGACTTGGGATTCAAGT

20

9

**F16M20TR IGF Arabidopsis thaliana g nomic cl ne F16M20, g nomic survey
s qu nc**

20

[illegible]

12

Tabl 10

F16A14-T7 IGF Arabidopsis thaliana genomic clone F16A14, genomic survey sequence

5 GGGGGGGATGGTACTGACTCCTATATGCCGAATCTTTGACATCTCTGTTTC
AATGGCCACAATCCTATTGAATCAGCTATATTAAGAAAATTATAACTCATCAA
ATAGCTTAAGACCATCGTTCCACGATCCTCACAATGCCTTNCNAGAGGAAC
TACCTTCCCGGAGTTAGTTCCCCATTCTGGGTTTACATCCATGAGACGGAAGA
10 GTAAGGTGACNATGGTCCATCGACGTGGATTGAATACNCTGTGGATCAGGAG
CTGTACGACCTGCTGGCTGATAAAGTAACCATGGCTTTAATCCTCCAAGAATA
TGAGCAACATATCCNAATGTAGACCTTGCACTTGTGACTATTTTATCAGTTAG
ACTTAGAAGATACNTCTCGGCGAGCGCCTTTTGGTCGTGTANCTTCTTGTCTT
NTGTTGAACCCTTTCTCCACTTGGCTGATNAACTTCAATGATCTCCCCTGCTG
15 AACTCGGTCGTTCCCAATACATGTTCTNTAAGGTNTCAGAGTACTCTGGATAC
NAAGATGTGACNAGAACAGCTNTAAGTGTCTGGCTTCTTGAATATATGACTTT
TGGCTCTTCTTGTGCACCTTGTTTCAAGGCAAAAGGTCTCTCTTCTGTCCAAC
TACAACTTGATCCNTTNCCTGTAAATTTCCCCCTCGAATGCTGAACTACCC
CTTCTCTAATAACCNCCCTCTCCTCCGCTCCTGAATAACTTCGGCTTGCTAGA
20 ATTCTCTCATTACCTCCCCACTTGAACCC CCCC GCGGTACAAACC

11

Table 11

T26M12-T7 TAMU Arabidopsis thaliana genomic clone T26M12, genomic
survey sequence [Arabidopsis thaliana]

5 ATTCGTGATGAGTACTATGCAAGCGAATCAAATGGTGA CTGCAGATACATTGT
ATGGCTAGCTAGGGACGGGCTTGGAACAGATTAACTACTCTTGCTTCCGTG
TTTCTCTACGCTATCTTGACTGAGAGAATCATTCTTGTTGACAACCGCAAGGA
TGTTAGTGATCTCTTATGTGAGCCATTTCCAGGTA CTTCATGGTTGCTTCCGC
10 TTGACTTTCCAATGCTGAATTATACTTATGCTTATGGCTACAATAAGGAATACC
TCGTTGTTACNGTACAATGTTGGAAAATCATGCCATCAACTCGACTTCAATTC
CGCCACATCTATATCTCCATAACATCCATGAATCTAGGGATANTGATAAGCTG
TTCTTCTGCCAAAANGGATCAAAGTTTTTTATCGACANATTTCCATGGGTAAAT
TAATTCANAACCAATGCCTTACTTTGGTTCCCAATCTTTATGGGCTGAAATCC
15 CANCTTTTCCAN ACCAAAACTAAGTTTAAGCTTATCCCCGGCAGAAAAGG

12

Tabl 12

ATTS3691 Gif-S dA Arabidopsis thaliana cDNA clone YAY241, mRNA
sequence [Arabidopsis thaliana]

5 AATGGTGATCTTGGGAATAGGATGCTGAGTCTAGCTTCAGCTTTTCTTTATGC
TCTCTTAACAAATAGGTTTTAACTTGTGGAAGTAGGAGTTGACATGGCTGACC
TTTTCTGCAAGCCATTTCAAACACTACTTGGTTTCTCCCCCAGAGTTTCCG
CTCAACAGCCACTTCAACGAGCAGTCTCTTCTACGCAATTCTGGCAACCCGA
10 TGGTTGCATATCGACATGTAGTTCGTGAATTCCAGTGACCAACAAAAGCTTTT
CTTTTGTGAGGATAGTCAAGTTTTGT

13,

**Tabl 13 T22N7TRB TAMU Arabidopsis thaliana g nomic cl n T22N7,
gen mic survey sequ nce [Arabid psis thaliana]**

5 CAAGCTTCGAGACAAGATATTCAGACGGCTTGTATGGTGAAGGCTTGCGATA
CAAGAAAGTTTTATGATACCTACTCAAGCAAGAACCTTCATCGAAATCTGCGG
TAAAAGCCCTCCTATTAGTCTATCCCGCCCCGGTGTTTCGGATTCTGTCTGAA
CATATAACAAAAAAAAAAGGTCAAAAGGAGAATTCTTTGAGCTAACAATG

3 4 5 6

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

Tabl 14

**31198 Lambda-PRL2 Arabidopsis thaliana cDNA clone 170K19XP 3', mRNA
s quenc [Arabidopsis thaliana]**

5 AAANNCCTTAANCAANTTTTACCGAANTCAAGGCGTTTACCCACTTCTCNCCN
GGTTTTAAGGTTCAAGGCGNNTTTTGGNAACCCNACAGTGATGGNGAGTTAT
CCGCGTTTACAANCCGACTACAAGGCTTCCAAAAACCCCGNGGAACNTGG
AANTTAAGAGANCATGGCTGAGATATACCTTCTGAGTTGTTCTGATGCNCTG
GTGGTCACAGGTTTATGGTCCTCACTCGTGGAGGTTGCCTCATGGCCTTGGA
10 GGGTTGAAGCCATGNGTGTTGAACAAAGCTGAGAATGGGACTGCCCATGAG
CCTTACTGTGTGAAAGCAAGATCAATAGAGCCCTGTTCCCAAGCGACATTGT
TCCATGGCTGTAAAGATTGAAACATGAATAGAGTCTCGAGGGCTTTTTTTGCC
TTAATAGATGTTGTACGGTCAAGAATTTTCAGAGTTGCCCAATAGACACGTAA
GGAATATTAGGATTAAGTATGTATCAGTTCATGACTTGATCGAGTTCTATATTC
15 TTTTCAAT

seq 1015